

On local invariants of pure three-qubit states

Anthony Sudbery

Department of Mathematics
University of York
Heslington
York
England YO1 5DD
Email: as2@york.ac.uk

24 February 2000, revised 15 November 2000

Abstract

We study invariants of three-qubit states under local unitary transformations, i.e. functions on the space of entanglement types, which is known to have dimension 6. We show that there is no set of six algebraically independent polynomial invariants of degree ≤ 6 , and find such a set with maximum degree 8. We describe an intrinsic definition of a canonical state on each orbit, and discuss the (non-polynomial) invariants associated with it.

1 Introduction

The invariants of many-particle states under unitary transformations which act on single particles separately (“local” transformations) are of interest [3, 7, 9, 10, 12] because they give the finest discrimination between different types of entanglement. They can be regarded as coordinates on the space of entanglement types (equivalently, the space of orbits of the group of local transformations). In this paper we study the case of pure states of three spin- $\frac{1}{2}$ particles, or qubits. For mixed states of two qubits, it is possible to give a complete set of invariants [11], describing the 9-dimensional space of orbits in terms of 18 invariants, nine of which may be taken to have only discrete values (for example, the signs of certain polynomials listed in [11]). For pure three-qubit states, where the space of orbits is known [3] to be 6-dimensional, we can at present do no more than find a set of six algebraically independent invariants. We will show (Section 3) that in order to do this with polynomials in the state coordinates it is necessary to go to polynomials of order 8, and we will exhibit (Section 4) a set of six independent invariants; their physical meaning is discussed in Section 5. We will also discuss (Section 6) the possibility of finding a more convenient set of non-polynomial invariants. Section 2 is an introductory discussion of the invariants of pure n -qubit states.

2 Pure states: general considerations

A general theory of local invariants of mixed n -particle states has been given by Rains and Grassl et al. [7, 12]. Here we review the part of that theory that refers to pure states.

The most general system is that of n non-identical particles A, B, \dots with one-particle state spaces of dimensions d_A, d_B, \dots . Let $\{|\psi_i^X\rangle : i = 1, \dots, d_X\}$ be an orthonormal basis of one-particle states of particle X ; then the general n -particle state can be written

$$|\Psi\rangle = \sum_{ijk\dots} t^{ijk\dots} |\psi_i^{(A)}\rangle |\psi_j^{(B)}\rangle |\psi_k^{(C)}\rangle \dots$$

where the sum is over values of i from 1 to d_A , values of j from 1 to d_B , and so on. By the First Fundamental Theorem of invariant theory [15] applied to $U(d_A), U(d_B), \dots$, any polynomial in $t^{ijk\dots}$ which is invariant under the action on $|\Psi\rangle$ of the local group $U(d_A) \times U(d_B) \times \dots$ is a sum of homogeneous polynomials of even degree (say $2r$), of the form

$$P_{\sigma\tau\dots}(\mathbf{t}) = t^{i_1 j_1 k_1 \dots} \dots t^{i_r j_r k_r \dots} \bar{t}_{i_1 j_{\sigma(1)} k_{\tau(1)} \dots} \dots \bar{t}_{i_r j_{\sigma(r)} k_{\tau(r)} \dots} \quad (2.1)$$

where σ, τ, \dots are permutations of $(1, \dots, r)$. Here $\bar{t}_{ijk\dots}$ is the complex conjugate of $t^{ijk\dots}$, and we adopt the usual summation convention on repeated indices, one in the upper position and one in the lower. Note that $P_{\sigma\tau\dots}$ is unchanged by simultaneous conjugation of the permutations σ, τ, \dots :

$$P_{\sigma\tau\dots}(\mathbf{t}) = P_{\sigma'\tau'\dots}(\mathbf{t}) \quad \text{if} \quad \sigma' = \kappa\sigma\kappa^{-1}, \quad \tau' = \kappa\tau\kappa^{-1}, \quad \dots$$

since such a conjugation merely expresses the effect of changing the order of the factors in each summand in P .

For two particles A, B there is just one permutation σ , which we can decompose into cycles $\kappa_1, \dots, \kappa_s$ of orders l_1, \dots, l_s with $l_1 + \dots + l_s = r$. The polynomial $P_\sigma(\mathbf{t})$ then splits into a product of polynomials $P_{\kappa_1} \cdots P_{\kappa_s}$, where P_κ depends only on the order of the cycle κ , which is equal to half the degree of P_κ :

$$\begin{aligned} P_\kappa(\mathbf{t}) &= t^{i_1 j_1} \bar{t}_{i_1 j_{\kappa(1)}} t^{i_{\kappa(1)} j_{\kappa(1)}} \bar{t}_{i_{\kappa(1)} j_{\kappa^2(1)}} \cdots \\ &= t^{i_1 j_1} \bar{t}_{i_1 j_2} t^{i_2 j_2} \bar{t}_{i_2 j_3} \cdots t^{i_l j_l} \bar{t}_{i_l j_1} \end{aligned}$$

(by renaming the dummy indices $j_{\kappa(1)}, j_{\kappa^2(1)}, \dots, j_{\kappa^{l-1}(1)}$)

$$= \text{tr}(\rho_B^l)$$

where $\rho_B = \text{tr}_A |\Psi\rangle\langle\Psi|$ is the density matrix of particle B , with matrix elements

$$(\rho_B)_k^j = t^{ij} \bar{t}_{ik}.$$

Thus the polynomial invariants of a two-particle pure state are the sums of the powers of the eigenvalues of ρ_B . These can all be expressed in terms of the first d_B power-sums, which generate the algebra of invariant polynomials and are algebraically independent if the eigenvalues are independent. However, they are not independent if $d_A < d_B$, for in that case some of the eigenvalues of ρ vanish. But clearly the same argument could be used to show that the algebra of invariants is generated by the traces of the powers of ρ_A , which is consistent because the non-zero eigenvalues of ρ_A are the same as those of ρ_B . Thus the algebra of polynomial invariants of two-particle pure states has a set of independent generators

$$\text{tr}(\rho_A^l) = \text{tr}(\rho_B^l), \quad l = 1, \dots, \min(d_A, d_B).$$

The non-zero eigenvalues of ρ_A (or ρ_B) are in fact the squares of the coefficients in the Schmidt decomposition of $|\Psi\rangle$, so what we have here is the well-known fact that the local invariants of a pure two-particle state are the symmetric functions of the Schmidt coefficients.

3 Polynomial invariants of three-qubit states

For the remainder of the paper we consider three spin- $\frac{1}{2}$ particles A, B, C . The classification of pure states of this system has been discussed in [4, 14], and their invariants in [5, 8]. It is known [9] that the dimension of the space of orbits is 6; there are therefore six algebraically independent local invariants. We will show that there are no more than five algebraically independent invariants of degree less than 8, and exhibit a set of six algebraically independent invariants with maximum degree 8.¹

The vector space of homogeneous invariants of degree $2r$ is spanned by functions $P_{\sigma\tau}$ labelled by pairs of elements of S_r , the group of permutations of r things. Thus there is one independent invariant of degree 2,

$$I_1 = P_{ee}(\mathbf{t}) = t^{ij}k\bar{t}_{ijk} = \langle \Psi | \Psi \rangle$$

where e is the identity permutation, so that $S_1 = \{e\}$. If $S_2 = \{e, \sigma\}$, the four linearly independent quartic invariants are

$$\begin{aligned} P_{ee}(\mathbf{t}) &= t^{i_1 j_1 k_1} \bar{t}_{i_1 j_1 k_1} t^{i_2 j_2 k_2} \bar{t}_{i_2 j_2 k_2} = \langle \Psi | \Psi \rangle^2, \\ I_2 = P_{e\sigma}(\mathbf{t}) &= t^{i_1 j_1 k_1} \bar{t}_{i_1 j_1 k_2} t^{i_2 j_2 k_2} \bar{t}_{i_2 j_2 k_1} = \text{tr}(\rho_C^2), \\ I_3 = P_{\sigma e}(\mathbf{t}) &= t^{i_1 j_1 k_1} \bar{t}_{i_1 j_2 k_1} t^{i_2 j_2 k_2} \bar{t}_{i_2 j_1 k_2} = \text{tr}(\rho_B^2), \\ I_4 = P_{\sigma\sigma}(\mathbf{t}) &= t^{i_1 j_1 k_1} \bar{t}_{i_1 j_2 k_2} t^{i_2 j_2 k_2} \bar{t}_{i_2 j_1 k_1} = \text{tr}(\rho_A^2) \end{aligned}$$

where ρ_A, ρ_B, ρ_C are the one-particle density matrices:

$$\rho_X = \text{tr}_{YZ} |\Psi\rangle\langle\Psi| \text{ where } \{X, Y, Z\} = \{A, B, C\} \text{ in some order.}$$

Thus there are at most four algebraically independent invariants of degree ≤ 4 .

Higher-order invariants $P_{\pi\sigma}(\mathbf{t})$ with $\pi, \sigma \in S_3$ are functions of the four quadratic and quartic invariants if π and σ are equal or if either of them is the identity. To see this, note first that if $\pi = \sigma$,

$$\begin{aligned} P_{\sigma\sigma}(\mathbf{t}) &= t^{i_1 j_1 k_1} \dots t^{i_r j_r k_r} \bar{t}_{i_1 j_{\sigma(1)} k_{\sigma(1)}} \dots \bar{t}_{i_r j_{\sigma(r)} k_{\sigma(r)}} \\ &= (\rho_A)_{i_{\tau(1)}}^{i_1} (\rho_A)_{i_{\tau(2)}}^{i_2} \dots (\rho_A)_{i_{\tau(r)}}^{i_r} \end{aligned}$$

where $\tau = \sigma^{-1}$. This is a product of traces of powers of ρ_A . But since ρ_A is a 2×2 matrix, the Cayley-Hamilton theorem enables us to express $\text{tr}(\rho_A^r)$ for $r \geq 3$ as a function of $\text{tr} \rho_A$ and $\text{tr} \rho_A^2$.

¹I understand that similar conclusions have been reached by Markus Grassl [6].

Secondly, if $\pi = e$,

$$\begin{aligned} P_{e\sigma}(\mathbf{t}) &= t^{i_1 j_1 k_1} \dots t^{i_r j_r k_r} \bar{t}_{i_1 j_1 k_{\sigma(1)}} \dots \bar{t}_{i_r j_r k_{\sigma(r)}} \\ &= (\rho_C)_{k_{\sigma(1)}}^{k_1} \dots (\rho_C)_{k_{\sigma(r)}}^{k_r} \end{aligned}$$

which is a product of traces of powers of ρ_C ; and similarly $P_{\pi e}(\mathbf{t})$ is a product of traces of powers of ρ_B .

Thus the only sextic invariants $P_{\pi\sigma}$ which might be algebraically independent of the quadratic and quartic invariants are those for which π and σ are distinct 2-cycles, or distinct 3-cycles, or one is a 2-cycle and the other is a 3-cycle. Moreover, in each of these categories all the possible pairs (π, σ) are related by simultaneous conjugation and therefore give the same invariant. There are therefore three possible independent sextic invariants:

1. π, σ distinct 3-cycles, say $\pi = (123), \sigma = (132)$. This gives

$$\begin{aligned} I_5 = P_{(123)(132)}(\mathbf{t}) &= t^{i_1 j_1 k_1} t^{i_2 j_2 k_2} t^{i_3 j_3 k_3} \bar{t}_{i_1 j_2 k_3} \bar{t}_{i_2 j_3 k_1} \bar{t}_{i_3 j_1 k_2} \\ &= (\rho_{BC})_{j_2 k_3}^{j_1 k_1} (\rho_{BC})_{j_3 k_1}^{j_2 k_2} (\rho_{BC})_{j_1 k_2}^{j_3 k_3} \end{aligned} \quad (3.1)$$

where $\rho_{BC} = \text{tr}_A |\Psi\rangle\langle\Psi|$ is the density matrix of the two-particle system of particles B and C . This invariant was identified by Kempe [8] as one which distinguishes three-particle states which have identical density matrices for every subsystem. It has exactly the same form when expressed as a function of ρ_{AB} or of ρ_{AC} .

2. π, σ distinct 2-cycles, say $\pi = (12), \sigma = (23)$. This gives

$$\begin{aligned} I'_5 = P_{(12)(23)}(\mathbf{t}) &= t^{i_1 j_1 k_1} t^{i_2 j_2 k_2} t^{i_3 j_3 k_3} \bar{t}_{i_1 j_2 k_1} \bar{t}_{i_2 j_1 k_3} \bar{t}_{i_3 j_3 k_2} \\ &= (\rho_B)_{j_2}^{j_1} (\rho_C)_{k_2}^{k_3} (\rho_{BC})_{j_1 k_3}^{j_2 k_2} \\ &= \text{tr}[(\rho_B \otimes \rho_C) \rho_{BC}]. \end{aligned} \quad (3.2)$$

3. π a 2-cycle, say (12) , and σ a 3-cycle, say (123) , or vice versa. These give

$$\begin{aligned} I''_5 = P_{(12)(123)}(\mathbf{t}) &= t^{i_1 j_1 k_1} t^{i_2 j_2 k_2} t^{i_3 j_3 k_3} \bar{t}_{i_1 j_2 k_2} \bar{t}_{i_2 j_1 k_3} \bar{t}_{i_3 j_3 k_1} \\ &= (\rho_{AC})_{i_2 k_3}^{i_1 k_1} (\rho_A)_{i_1}^{i_2} (\rho_C)_{k_1}^{k_3} \\ &= \text{tr}[(\rho_A \otimes \rho_C) \rho_{AC}] \end{aligned} \quad (3.3)$$

and

$$\begin{aligned} I'''_5 = P_{(123)(12)}(\mathbf{t}) &= t^{i_1 j_1 k_1} t^{i_2 j_2 k_2} t^{i_3 j_3 k_3} \bar{t}_{i_1 j_2 k_2} \bar{t}_{i_2 j_3 k_1} \bar{t}_{i_3 j_1 k_3} \\ &= \text{tr}[(\rho_A \otimes \rho_B) \rho_{AB}]. \end{aligned} \quad (3.4)$$

Primes have been placed on the symbols for these last three invariants because they will not feature in our final list of independent invariants, each of them being expressible in terms of I_5 and the quadratic and quartic invariants. To show this, we write I_5 in terms of 2×2 matrices by considering the 4×4 matrix ρ_{BC} as a set of four 2×2 matrices $X_{j_2}^{j_1}$: the matrix elements of $X_{j_2}^{j_1}$, labelled by (k_1, k_2) , are

$$(X_{j_2}^{j_1})_{k_2}^{k_1} = (\rho_{BC})_{j_2 k_2}^{j_1 k_1}.$$

Then

$$I_5 = \text{tr}(X_{j_2}^{j_1} X_{j_1}^{j_3} X_{j_3}^{j_2}).$$

Now we use the 2×2 matrix identity

$$\begin{aligned} \text{tr}(XYZ) + \text{tr}(XZY) &= \text{tr} X \text{tr}(YZ) + \text{tr} Y \text{tr}(ZX) + \text{tr} Z \text{tr}(XY) \\ &\quad - \text{tr} X \text{tr} Y \text{tr} Z \end{aligned} \quad (3.5)$$

which holds for any 2×2 matrices X, Y, Z , and can be obtained by trilinearising (or ‘‘polarising’’ [15] — replace X first by $X + Y$ and then by $X + Y + Z$) the cubic identity

$$\text{tr} X^3 = \frac{3}{2} \text{tr} X \text{tr} X^2 - \frac{1}{2} (\text{tr} X)^3$$

which in turn is obtained by taking the trace of the Cayley-Hamilton theorem. Apply (3.5) to the matrices $X_{j_2}^{j_1}, X_{j_1}^{j_3}, X_{j_3}^{j_2}$ occurring in the expression for I_5 . The first term on the left-hand side is I_5 ; the second is

$$\text{tr}(X_{j_2}^{j_1} X_{j_3}^{j_2} X_{j_1}^{j_3}) = \text{tr}(\rho_{BC}^3) = \text{tr}(\rho_A^3)$$

since the non-zero eigenvalues of ρ_{BC} are the same as those of ρ_A (both being the squares of the coefficients in a Schmidt decomposition of $|\Psi\rangle$). The first term on the right-hand side is

$$\begin{aligned} \text{tr}(X_{j_2}^{j_1}) \text{tr}(X_{j_1}^{j_3} X_{j_3}^{j_2}) &= (\rho_B)_{j_2}^{j_1} (\rho_{BC})_{j_1 k_2}^{j_3 k_1} (\rho_{BC})_{j_3 k_1}^{j_2 k_2} \\ &= (\rho_B)_{j_2}^{j_1} t^{i_1 j_3 k_1} \bar{t}_{i_1 j_1 k_2} t^{i_2 j_2 k_2} \bar{t}_{i_2 j_3 k_1} \\ &= (\rho_B)_{j_2}^{j_1} (\rho_A)_{i_2}^{i_1} (\rho_{AB})_{i_1 j_1}^{i_2 j_2} \\ &= \text{tr}[(\rho_A \otimes \rho_B) \rho_{AB}]; \end{aligned}$$

the second and third terms differ from the first only by permuting the indices j_1, j_2, j_3 and therefore (after summing) are equal to it; and the last term is

$$\text{tr}(X_{j_2}^{j_1}) \text{tr}(X_{j_3}^{j_2}) \text{tr}(X_{j_1}^{j_3}) = \text{tr}(\rho_B^3).$$

Thus (3.5) gives

$$I_5 = 3 \operatorname{tr}[(\rho_A \otimes \rho_B)\rho_{AB}] - \operatorname{tr}(\rho_A^3) - \operatorname{tr}(\rho_B^3). \quad (3.6)$$

Similarly, using the alternative expressions for I_5 in terms of ρ_{AB} and ρ_{AC} gives

$$I_5 = 3 \operatorname{tr}[(\rho_B \otimes \rho_C)\rho_{BC}] - \operatorname{tr}(\rho_B^3) - \operatorname{tr}(\rho_C^3) \quad (3.7)$$

$$= 3 \operatorname{tr}[(\rho_A \otimes \rho_C)\rho_{AC}] - \operatorname{tr}(\rho_A^3) - \operatorname{tr}(\rho_C^3). \quad (3.8)$$

So there are at most five independent invariants of degree 6 or less. Since six invariants are needed to parametrise the orbits [9], we must use at least one invariant of degree 8 or more. A convenient, and physically significant, choice is the 3-tangle identified by Coffman, Kundu and Wootters [5]:

$$I_6 = \frac{1}{4} \mathcal{T}_{123}^2 = \left| \epsilon_{i_1 i_2} \epsilon_{i_3 i_4} \epsilon_{j_1 j_2} \epsilon_{j_3 j_4} \epsilon_{k_1 k_3} \epsilon_{k_2 k_4} t^{i_1 j_1 k_1} t^{i_2 j_2 k_2} t^{i_3 j_3 k_3} t^{i_4 j_4 k_4} \right|^2 \quad (3.9)$$

where ϵ_{ij} is the antisymmetric tensor in two dimensions ($\epsilon_{12} = -\epsilon_{21} = 1$, $\epsilon_{11} = \epsilon_{22} = 0$). The expression between the modulus signs is an $SU(2)^3$ invariant (though not a $U(2)^3$ invariant — its phase is not invariant under local transformations), so its modulus is a local invariant. The invariant I_6 can be put into our standard form of a sum of terms like (2.1) by multiplying the $SU(2)^3$ invariant by its complex conjugate

$$\epsilon^{i_5 i_6} \epsilon^{i_7 i_8} \epsilon^{j_5 j_6} \epsilon^{j_7 j_8} \epsilon^{k_5 k_7} \epsilon^{k_6 k_8} \bar{t}_{i_5 j_5 k_5} \bar{t}_{i_6 j_6 k_6} \bar{t}_{i_7 j_7 k_7} \bar{t}_{i_8 j_8 k_8}$$

(where the contravariant tensor ϵ^{ij} is numerically the same as ϵ_{ij}), and using the identity

$$\epsilon^{ab} \epsilon_{cd} = \delta_c^a \delta_d^b - \delta_d^a \delta_c^b.$$

To show that the invariants I_1, \dots, I_6 are independent it is sufficient to show that their gradients are linearly independent at some point. To calculate these gradients in the 16-(real)dimensional space of pure states, we can treat t^{ijk} and \bar{t}_{ijk} formally as independent coordinates; the fact that our invariants are real means that the 16 components of the gradient of I_a are the real and imaginary parts of the partial derivatives with respect to t^{ijk} . The results of calculating $\partial I_a / \partial t^{ijk}$ and putting

$$t^{000} = t^{010} = t^{110} = 0, \quad t^{011} = t^{100} = t^{101} = t^{111} = 1, \quad t^{001} = i,$$

$$\bar{t}^{ijk} = \text{complex conjugate of } t^{ijk}$$

(where 0 and 1 are the two possible values of i, j, k) are as follows:

$$\begin{aligned}
\partial_t I_1 &= (0, -i, 0, 1, 1, 1, 0, 1) \\
\partial_t I_2 &= (-2i, -8i, 2, 8, 4, 10, 2, 8) \\
\partial_t I_3 &= (0, 2 - 8i, 0, 6 - 2i, 6, 8 - 2i, 2 + 2i, 6 + 2i) \\
\partial_t I_4 &= (2 - 2i, 2 - 6i, 0, 6 - 2i, 6, 8 - 2i, 0, 8 + 2i) \\
\partial_t I_5 &= (6 - 9i, 12 - 36i, 6, 30 - 12i, 21, 45 - 12i, 9 + 6i, 36 + 12i) \\
\partial_t I_6 &= (-8, 0, -8 + 16i, 8, 8, 0, -8i, 0)
\end{aligned}$$

These six vectors are indeed linearly independent over \mathbb{R} .

4 Physical significance of the invariants

The invariant I_1 is just the norm of the three-party state and therefore has no physical significance; we will normally set it equal to 1. The three invariants I_1, I_2, I_3 are one-particle quantities, giving the eigenvalues of the one-particle density matrices; they are equivalent to the one-particle entropies, which measure how entangled each particle is with the other two together. The entanglement in each pair of particles and the three-way entanglement of the whole system are all given by the last invariant I_6 , as follows. A good measure of the entanglement of two qubits A, B in a mixed state is the *2-tangle* τ_{AB} [5], which is a monotonic function of the entanglement of formation [16]. The three-way entanglement of three qubits A, B, C in a pure state is measured by the *3-tangle*

$$\tau_{ABC} = \tau_{A(BC)} - \tau_{AB} - \tau_{AC}$$

where $\tau_{A(BC)} = 4 \det \rho_A = 2(I_1^2 - I_2)$ is another measure (equivalent to the entropy of A) of how entangled A is with the pair (BC) . It can be shown [5] that τ_{ABC} is invariant under permutations of A, B and C ; in fact it is equal to our invariant I_6 . By solving the equations expressing the permutation invariance of τ_{ABC} , we can now give formulae for all three 2-tangles and the 3-tangles in terms of our invariants:

$$\begin{aligned}
\tau_{AB} &= 1 - I_2 - I_3 + I_4 - \frac{1}{2}I_6, \\
\tau_{AC} &= 1 - I_2 + I_3 - I_4 - \frac{1}{2}I_6, \\
\tau_{BC} &= 1 + I_2 - I_3 - I_4 - \frac{1}{2}I_6, \\
\tau_{ABC} &= I_6.
\end{aligned}$$

The 3-tangle I_6 is maximal for the GHZ state $|000\rangle + |111\rangle$, whose 2-tangles vanish; on the other hand, I_6 vanishes at the states $p|100\rangle + q|010\rangle + r|001\rangle$.

The remaining invariant, I_5 , is a different and independent measure of the entanglement of each pair of qubits. Its existence shows that the 2-tangles and 3-tangle are not sufficient to determine a pure 3-qubit state up to local equivalence. As is shown by eqs. (3.6)–(3.8), this invariant is equivalent to any one of the two-qubit quantities $\kappa_{AB} = \text{tr}[(\rho_A \otimes \rho_B)\rho_{AB}]$ (together with one-qubit quantities), and it relates these three 2-qubit quantities to each other. If we regard the hermitian operators ρ_A and ρ_B as observables, then κ_{AB} is the expectation value of $\rho_A\rho_B$, so $\kappa_{AB} - I_2I_3$ is the correlation between the eigenvalues of ρ_A and ρ_B . It is related to the relative entropy of the two-qubit state ρ_{AB} relative to the product state $\rho_A \otimes \rho_B$, and is a second measure of the entanglement of the pair (A, B) , independent of the 2-tangle τ_{AB} .

Finally, we give the values of these invariants for some special states (all of which are taken to be normalised).

For a factorised state $a|111\rangle + b|100\rangle$,

$$I_2 = 1, \quad I_3 = I_4 = a^4 + b^4, \quad I_5 = a^6 + b^6, \quad I_6 = 0.$$

For a generalised GHZ state $p|000\rangle + q|111\rangle$,

$$I_2 = I_3 = I_4 = p^4 + q^4, \quad I_5 = p^6 + q^6, \quad I_6 = 4p^2q^2.$$

For the minimally 3-tangled [5] state $p|100\rangle + q|010\rangle + r|001\rangle$,

$$I_2 = p^4 + (q^2 + r^2)^2, \quad I_3 = q^4 + (r^2 + p^2)^2, \quad I_4 = r^4 + (p^2 + q^2)^2, \\ I_5 = p^6 + q^6 + r^6 + 3p^2q^2r^2, \quad I_6 = 0.$$

5 Canonical coordinates

An alternative type of invariant, not necessarily a polynomial in the coordinates of the state vector, is obtained by specifying a canonical point on each orbit. The values of the invariant functions at any point are then the coordinates of the canonical point on its orbit. The canonical points lie on a manifold corresponding to the space of orbits, and their coordinates can (at least locally) be expressed in terms of an appropriate number of parameters.

One form of canonical state was suggested independently by Linden and Popescu [9] and by Schlienz [13], who pointed out that any pure state of three qubits can be written as

$$|\Psi\rangle = \cos\theta|0\rangle(\cos\phi|0\rangle|0\rangle + \sin\phi|1\rangle|1\rangle) \\ + \sin\theta|1\rangle(r(-\sin\phi|0\rangle|0\rangle + \cos\phi|1\rangle|1\rangle) + s|0\rangle|1\rangle + te^{i\omega}|1\rangle|0\rangle) \quad (5.1)$$

where $0 \leq \theta, \phi \leq \pi/4$, $0 \leq \omega < 2\pi$, and r, s, t are non-negative real numbers satisfying $r^2 + s^2 + t^2 = 1$. Simpler canonical forms, in which the number of non-zero coefficients is reduced to five, have since been proposed by Acin et al [1] and Carteret et al [2]; the latter form is

$$p|100\rangle + q|010\rangle + r|001\rangle + s|111\rangle + te^{i\theta}|000\rangle$$

where p, q, r, s, t and θ are real parameters. It is straightforward to calculate the invariants I_1, \dots, I_6 in terms of either of the above sets of parameters; the results are not enlightening.

We will now describe another, more intrinsically defined, form of canonical point whose coordinates are more simply related to I_1, \dots, I_6 .

The three-particle state $|\Psi\rangle$ has three Schmidt decompositions:

$$\begin{aligned} |\Psi\rangle &= \sum_i \alpha_i |\phi_i\rangle_A |\Phi_i\rangle_{BC} \\ &= \sum_i \beta_i |\theta_i\rangle_B |\Theta_i\rangle_{AC} \\ &= \sum_i \gamma_i |\chi_i\rangle_C |X_i\rangle_{AB} \end{aligned} \tag{5.2}$$

where $\{|\phi_i\rangle\}$, $\{|\theta_i\rangle\}$ and $\{|\chi_i\rangle\}$ ($i = 0, 1$) are orthonormal pairs of one-particle states, $\{|\Phi_i\rangle\}$, $\{|\Theta_i\rangle\}$ and $\{|X_i\rangle\}$ are orthonormal pairs of two-particle states, the suffices indicate which of the three particles A, B, C are in which state, and $\{\alpha_i\}$, $\{\beta_i\}$ and $\{\gamma_i\}$ are pairs of non-negative real numbers satisfying

$$\alpha_1^2 + \alpha_2^2 = \beta_1^2 + \beta_2^2 = \gamma_1^2 + \gamma_2^2 = \langle \Psi | \Psi \rangle = I_1. \tag{5.3}$$

These Schmidt coefficients, being the positive square roots of the eigenvalues of the one-particle density matrices ρ_A, ρ_B, ρ_C , are related to the quartic invariants by

$$\begin{aligned} \alpha_1^4 + \alpha_2^4 &= \text{tr}(\rho_A^2) = I_2, \\ \beta_1^4 + \beta_2^4 &= \text{tr}(\rho_B^2) = I_3, \\ \gamma_1^4 + \gamma_2^4 &= \text{tr}(\rho_C^2) = I_4. \end{aligned} \tag{5.4}$$

These equations have unique real non-negative solutions for $\alpha_i, \beta_i, \gamma_i$ provided the invariants I_1, \dots, I_4 satisfy

$$I_1 > 0, \quad \frac{1}{2}I_1^2 \leq I_2, I_3, I_4 \leq I_1^2.$$

Now consider the coordinates c^{ijk} of $|\Psi\rangle$ with respect to the canonical basis $|\phi_i\rangle_A |\theta_j\rangle_B |\chi_k\rangle_C$. If the states $|\phi_i\rangle, |\theta_j\rangle, |\chi_k\rangle$ were uniquely determined by $|\Psi\rangle$

— and they almost are — then the coordinates c^{ijk} would be local invariants. However, the Schmidt decompositions do not determine the phases of $|\phi_i\rangle$, $|\theta_j\rangle$ and $|\chi_k\rangle$. We can fix these by requiring that four of the c^{ijk} should be real: for example, we can change the phases of $|\phi_0\rangle$ and $|\phi_1\rangle$ to make c^{000} and c^{100} real, then change the phases of $|\theta_0\rangle$ and $|\theta_1\rangle$ to make c^{001} and c^{011} real, simultaneously changing the phase of $|\chi_0\rangle$ to keep c^{000} and c^{100} real. (It is easy to show that under the six-dimensional group of phase changes of the basis vectors, the generic set of coordinates has two-dimensional stabiliser, so that the orbits are four-dimensional and therefore four phases can be removed.)

From the Schmidt decompositions we obtain the one-particle density matrices

$$\begin{aligned}\rho_A &= \sum_i \alpha_i^2 |\phi_i\rangle\langle\phi_i|, \\ \rho_B &= \sum_i \beta_i^2 |\theta_i\rangle\langle\theta_i|, \\ \rho_C &= \sum_i \gamma_i^2 |\chi_i\rangle\langle\chi_i|.\end{aligned}\tag{5.5}$$

Hence the coordinates c^{ijk} satisfy

$$\begin{aligned}\sum_{jk} c^{ijk} \bar{c}_{ljk} &= \alpha_i^2 \delta_l^i, \\ \sum_{ik} c^{ijk} \bar{c}_{imk} &= \beta_j^2 \delta_m^j, \\ \sum_{ij} c^{ijk} \bar{c}_{ijn} &= \gamma_k^2 \delta_n^k.\end{aligned}\tag{5.6}$$

To obtain a relation between the c^{ijk} and Kempe's invariant I_5 , we calculate

$$\begin{aligned}& \text{tr}[(\rho_A \otimes \rho_B) \rho_{AB}] \\ &= \text{tr} \left[\left(\sum_i \alpha_i^2 |\phi_i\rangle_A \langle\phi_i|_A \right) \left(\sum_j \beta_j^2 |\theta_j\rangle_B \langle\theta_j|_B \right) \left(\sum_k \gamma_k^2 |X_k\rangle_{AB} \langle X_k|_{AB} \right) \right] \\ &= \sum_{ijk} \alpha_i^2 \beta_j^2 \gamma_k^2 |\langle\phi_i|\theta_j|X_k\rangle|^2.\end{aligned}$$

But

$$c^{ijk} = \langle\phi_i|_A \langle\theta_j|_B \langle\chi_k|_C | \Psi \rangle = \gamma_k \langle\phi_i|_A \langle\theta_j|_B | X_k \rangle_{AB}.$$

Hence

$$\text{tr}(\rho_A \rho_B \rho_{AB}) = \sum_{ijk} \alpha_i^2 \beta_j^2 |c^{ijk}|^2$$

and so, using (3.6),

$$I_5 = 3 \sum_{ijk} \alpha_i^2 \beta_j^2 |c^{ijk}|^2 - \sum_i \alpha_i^6 - \sum_j \beta_j^6. \quad (5.7)$$

Finally, the relation between the c^{ijk} and the 3-tangle I_6 needs a longer argument which we will not give here. The result is

$$I_6 = \det R \quad (5.8)$$

where

$$R_j^i = (\alpha_i^4 + \alpha_i^2) \delta_j^i - \sum_{kl} (\beta_k^2 + \gamma_l^2) c^{ikl} \bar{c}_{jkl}$$

In order to determine how many states have the same values of the invariants I_1, \dots, I_6 , and therefore how many further discrete-valued invariants are needed to specify uniquely a pure state of three qubits up to local transformations, one would need to find the number of different sets of coordinates c^{ijk} satisfying the reality conditions given above and the equations (5.6), (5.7) and (5.8), where α_i, β_i and γ_i are determined by (5.3) and (5.4).

Acknowledgements

I am grateful to Bob Gingrich for finding an error in an earlier version of this paper and for performing the calculations whose result is reported at the end of Section 3.

References

- [1] A. Acin, A. Andrianov, L. Costa, E. Jane, J. I. Latorre, and R. Tarrach. Generalized Schmidt decomposition and classification of three-qubit states. *Phys. Rev. Lett.*, 85:1560, 2000. quant-ph/0003050.
- [2] H. A. Carteret, A. Higuchi, and A. Sudbery. Multipartite generalisation of the Schmidt decomposition. *J. Math. Phys.*, 2000. (to be published). quant-ph/0006125.

- [3] H. A. Carteret, N. Linden, S. Popescu, and A. Sudbery. Multi-particle entanglement. *Found. Phys.*, 29:527, 1999.
- [4] H. A. Carteret and A. Sudbery. Local symmetry properties of pure states of three qubits. *J. Phys. A*, 33:4981, 2000. quant-ph/0001091.
- [5] V. Coffman, J. Kundu, and W. K. Wootters. Distributed entanglement. *Phys. Rev. A*, 61(2306), 2000. quant-ph/9907047.
- [6] M. Grassl. talk presented at the Isaac Newton Institute, Cambridge in the workshop on complexity, computation and the physics of information, July 1999.
- [7] M. Grassl, M. Rötteler, and T. Beth. Computing local invariants of qubit systems. *Phys. Rev. A*, 58:1853, 1998. quant-ph/9712040.
- [8] J. Kempe. Multi-particle entanglement and its applications to cryptography. *Phys. Rev. A*, 60:910, 1999. quant-ph/9902036.
- [9] N. Linden and S. Popescu. On multi-particle entanglement. *Fortschr. Phys.*, 46:567, 1998. quant-ph/9711016.
- [10] N. Linden, S. Popescu, and A. Sudbery. Non-local parameters for multi-particle density matrices. *Phys. Rev. Lett.*, 83:243, 1999. quant-ph/9801076.
- [11] Y. Makhlin. Nonlocal properties of two-qubit gates and mixed states and optimization of quantum computations. 1999. quant-ph/0002045.
- [12] E. M. Rains. Polynomial invariants of quantum codes. *IEEE Trans. Information Theory*, 46:54, 2000. quant-ph/9704042.
- [13] J. Schlienz. PhD thesis.
- [14] J. Schlienz and G. Mahler. The maximal entangled three-particle state is unique. *Phys. Lett. A*, 224:39, 1996.
- [15] H. Weyl. *The Classical Groups*. Princeton, 1946.
- [16] W. K. Wootters. Entanglement of formation of an arbitrary state of two qubits. *Phys. Rev. Lett.*, 80:2245, 1998. quant-ph/9709029.