

MIT OpenCourseWare  
<http://ocw.mit.edu>

8.323 Relativistic Quantum Field Theory I  
Spring 2008

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.

8.323: Relativistic Quantum Field Theory I

Prof. Alan Guth

March 13, 2008

INFORMAL NOTES  
DIRAC DELTA FUNCTION AS A DISTRIBUTION

Why the Dirac Delta Function is not a Function:

The Dirac delta function  $\delta(x)$  is often described by considering a function that has a narrow peak at  $x = 0$ , with unit total area under the peak. In the limit as the peak becomes infinitely narrow, keeping fixed the area under the peak, the function is sometimes said to approach a Dirac delta function. One example of such a limit is


$$g(x) \equiv \lim_{\sigma \rightarrow 0} g_{\sigma}(x) , \quad (4.1)$$

where

$$g_{\sigma}(x) \equiv \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2}x^2/\sigma^2} . \quad (4.2)$$

The area under  $g_{\sigma}(x)$  is 1, for any value of  $\sigma > 0$ , and  $g_{\sigma}(x)$  approaches 0 as  $\sigma \rightarrow 0$  for any  $x$  other than  $x = 0$ .

However, it was pointed out long ago that the delta function cannot be rigorously defined this way. The function  $g(x)$  is equal to zero for any  $x \neq 0$ , and is infinite at  $x = 0$ ; it can be shown that any such function integrates to zero. To see this, define the integral as the area under the curve, and consider the construction:



In this picture the vertical axis is entirely encased in rectangles, each of which has height 1. The width of the rectangles vary, with the lowest rectangle having width  $\epsilon$ , for some  $\epsilon > 0$ , and each successive rectangle has half the width of the rectangle below. Note that the outline of the boxes is everywhere above the curve  $g(x)$ , so

the area under  $g(x)$  must be less than the total area of the boxes. But the total area of the boxes is given by a geometric series,

$$A = \epsilon + \frac{1}{2}\epsilon + \frac{1}{4}\epsilon + \dots \leq 2\epsilon . \quad (4.3)$$

Since  $\epsilon$  can be chosen as small as one likes, the area under the limit function  $g(x)$  must be zero.

This result does not contradict the statement that the area under  $g_\sigma(x)$  is 1 for any  $\sigma > 0$ . Rather, this is a case where the limit of an integral is not the same as the integral of the limit of the integrand. The integral has the value 1 for every  $\sigma > 0$ , so the limit of the integral as  $\sigma \rightarrow 0$  is 1. However, if one takes the limit of the integrand first, and then integrates, the answer is zero.

### Dirac Delta Function as a Distribution:

A Dirac delta function is defined to have the property that

$$\int_{-\infty}^{\infty} dx \varphi(x) \delta(x - a) \equiv \varphi(a) . \quad (4.4)$$

But we have just seen that there is no function  $\delta(x)$  which has this property, as long as integration is defined by the area under a curve. However, there is no problem defining a distribution that behaves this way. Using the notation we defined earlier for distributions, we define

$$T_{\delta(x-a)}[\varphi] \equiv \varphi(a) . \quad (4.5)$$

Here  $T_{\delta(x-a)}$  is the name that we will use for the distribution that acts like integration over the delta function  $\delta(x - a)$ , and  $\varphi(x)$  is the test function. This is certainly a linear functional that is defined on all Schwartz functions, and therefore is a tempered distribution. We interpret Eqs. (4.4) and (4.5) as meaning the same thing, where the second form emphasizes the definition as a distribution, and the first form emphasizes that distributions can be viewed as a generalized kind of function, with a notation that makes them look like functions. That is, we define

$$\int_{-\infty}^{\infty} dx \varphi(x) \delta(x - a) \equiv T_{\delta(x-a)}[\varphi] \equiv \varphi(a) . \quad (4.6)$$

To put it differently, we must remember that an integral over a delta function, such as Eq. (4.4), is not defined as a standard integral— instead it is symbolic integral, which is defined as a distribution which maps the function that multiplies the delta function to its value at the point where the argument of the delta function vanishes.

**What about  $\int dp e^{-ip(x-a)}$  ?**

If we interpret this integral in the sense of Riemann or Lebesgue, it simply does not exist — it diverges. As a distribution, however, we can interpret

$$I(x) \equiv \int_{-\infty}^{\infty} dp e^{-ip(x-a)} \quad (4.7)$$

as the Fourier transform of

$$f(p) = e^{ipa} . \quad (4.8)$$

Since  $I(x)$  is the Fourier transform of the distribution  $T_f[\varphi]$ , it must itself be a well-defined distribution. Using the notation of distributions, the distribution corresponding to  $I$  is then given by

$$T_I[\varphi] = \tilde{T}_f[\varphi] . \quad (4.9)$$

To evaluate the distribution  $T_I[\varphi]$ , we first write the distribution corresponding to function  $f(p)$  in the usual way:

$$T_f[\varphi] = \int_{-\infty}^{\infty} dp f(p) \varphi(p) = \int_{-\infty}^{\infty} dp e^{ipa} \varphi(p) . \quad (4.10)$$

The Fourier transform of this distribution is then defined by applying the same distribution to the Fourier transform of the test function, so

$$\tilde{T}_f[\varphi] \equiv T_f[\tilde{\varphi}] = \int_{-\infty}^{\infty} dp e^{ipa} \tilde{\varphi}(p) . \quad (4.11)$$

But the inverse Fourier transform is given by

$$\varphi(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} dp e^{ipx} \tilde{\varphi}(p) , \quad (4.12)$$

so by comparing the two formulas above, one sees that

$$\tilde{T}_f[\varphi] = 2\pi\varphi(a) . \quad (4.13)$$

But this is exactly the definition of the delta function distribution, as given in Eq. (4.6), so

$$\tilde{T}_f[\varphi] = 2\pi T_{\delta(x-a)}[\varphi] . \quad (4.14)$$

If we write these distributions as symbolic integrals, then the above equation becomes

$$\int_{-\infty}^{\infty} dx I(x)\varphi(x) = 2\pi \int_{-\infty}^{\infty} dx \delta(x-a)\varphi(x) , \quad (4.15)$$

where we remembered from Eq. (4.9) that  $\tilde{T}_f[\varphi]$  is equal to the distribution corresponding to  $I(x)$ . Since Eq. (4.15) holds for an arbitrary test function  $\varphi(x)$ , we can equate the distributions without showing their arguments:

$$I(x) = \int_{-\infty}^{\infty} dp e^{-ip(x-a)} = 2\pi\delta(x-a) . \quad (4.16)$$

### The Derivative of a Delta Function:

If a Dirac delta function is a distribution, then the derivative of a Dirac delta function is, not surprisingly, the derivative of a distribution. We have not yet defined the derivative of a distribution, but it is defined in the obvious way. We first consider a distribution corresponding to a function, and ask what would be the distribution corresponding to the derivative of the function.

Starting with a well-behaved (i.e., piecewise continuous and bounded by some power of  $t$ ) function  $f(t)$ , we defined the corresponding distribution by

$$T_f[\varphi] \equiv \int_{-\infty}^{\infty} dt f(t)\varphi(t) . \quad (4.17)$$

Then if we write the distribution corresponding to  $df/dt$ , we get

$$T_{df/dt}[\varphi] = \int_{-\infty}^{\infty} dt \frac{df}{dt} \varphi(t) \quad (4.18)$$

Since  $f(t)$  is bounded for large  $|t|$  by a power of  $t$ , and  $\varphi(t)$  falls off faster than any power, we can integrate by parts without encountering a surface term:

$$T_{df/dt}[\varphi] = - \int_{-\infty}^{\infty} dt f(t) \frac{d\varphi}{dt} = -T_f \left[ \frac{d\varphi}{dt} \right] . \quad (4.19)$$

This result can then be taken as the general definition of the derivative of a distribution:

$$T'[\varphi] \equiv -T \left[ \frac{d\varphi}{dt} \right] . \quad (4.20)$$

Applying this result to a delta function in the notation of Eq. (4.6), it looks exactly like integration by parts:

$$\int_{-\infty}^{\infty} dt f(t) \frac{d\delta(t-a)}{dt} = - \left. \frac{df}{dt} \right|_{t=a} . \quad (4.21)$$